

TECHNICAL DATA SHEET

CLUBZ HOSPITAL QUAT-CLEANER

DESCRIPTION

- **Clubz** is designed specifically as a general ready-to-use cleaner and disinfectant for use in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, shower rooms, locker rooms, public restrooms, schools, food processing plants, food service establishments, restaurants, commercial kitchens, office buildings, lodging establishments, retail businesses and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. For plastic and painted surfaces, spot test on an inconspicuous area before use. May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact with food.
- Clubz deodorizes those areas that generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms. It doesn't contain phosphates or phosphorous.
- When used as directed, Clubz is highly effective against a broad-spectrum of pathogenic microorganisms including bacteria, antibiotic resistant bacteria, viruses, fungi, mold and mildew.
- Clubz has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

FEATURES

- Kills 99.9% of bacteria
- Kills 99.9% of germs
- Disinfectant, cleaner, tuberculocidal
- Cuts grease and grime
- Eliminates odors
- Kills Pandemic 2009 H1N1 influenza A virus
- Meets OSHA Bloodborne Pathogen standard
- For HIV, HBV and HCV
- Doesn't contain phosphates or phosphorous



HOW TO USE

- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- This product is not for use on critical or semi-critical medical device surfaces.
- Fill or refill the 16 oz. CLUBZ HOSPITAL QUAT-CLEANER spray bottle with CLUBZ HOSPITAL QUAT-CLEANER refill. Pour product into spray bottle over sink or tub basin.
- DISINFECTION, DEODORIZING AND CLEANING Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 1 minute). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.
- TO DISINFECT FOOD PROCESSING PREMISES: Before using CLUBZ HOSPITAL QUAT-CLEANER, food products and packaging materials must be removed or carefully protected. For floors, walls and storage areas, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow CLUBZ HOSPITAL QUAT-CLEANER to penetrate and remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 1 minute). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be precleaned before using this product.
- TO DISINFECT FOOD SERVICE ESTABLISHMENT FOOD CONTACT SURFACES: Before using CLUBZ HOSPITAL QUAT-CLEANER food products and packaging materials must be removed or carefully protected. For countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow CLUBZ HOSPITAL QUAT-CLEANER to penetrate and remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 1 minute). Wipe off with a clean cloth, mop or sponge and thoroughly rine all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

PERSONAL PROTECTION

• When handling items soiled with blood or bodily fluids use disposable latex gloves, gowns, masks and eye coverings.

CLEANING PROCEDURES

• Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

CONTACT TIME

• Allow surface to remain wet for HBV and HVC 1 minute, for HIV-1 1 minute. These contact times will not control other common types of viruses and bacteria.



TECHNICAL DATA SHEET

STORAGE & DISPOSAL

- **PESTICIDE STORAGE** Store in a dry place no lower in temperature than 50°F or higher than 120°F.
- **PESTICIDE DISPOSAL** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
- **CONTAINER DISPOSAL** Nonrefillable container. Do not reuse or refill container. Clean container promptly after emptying.
- **Triple rinse as follows**: Fill container ¼ full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Offer for recycling or reconditioning, if available. If not available, puncture and dispose in a sanitary landfill.
- **Container disposal for Refill Spray Bottle:** Do not reuse empty spray bottle except with refill bottle, otherwise discard in trash. Do not reuse empty refill bottle. Wrap refill bottle and discard in trash. Offer for recycling or reconditioning, if available.